arXiv:1401.2836v1 [math.AC] 13 Jan 2014

NOTES ON ADDITIVELY DIVISIBLE COMMUTATIVE SEMIRINGS

TOMÁŠ KEPKA AND MIROSLAV KORBELÁŘ

Abstract. Commutative semirings with divisible additive semigroup are studied. We show that an additively divisible commutative semiring is idempotent, provided that it is finitely generated and torsion. In case that a one-generated additively divisible semiring posseses no unit, it must contain an ideal of idempotent elements. We also present a series of open questions about finitely generated commutative semirings and their equivalent versions.

It is well known that a commutative field is finite provided that it is a finitely generated ring. Consequently, no finitely generated commutative ring (whether unitary or not) contains a copy of the field $\mathbb{Q}$ of rational numbers. On the other hand, it seems to be an open problem whether a finitely generated (commutative) semiring S can contain a copy of the semiring (parasemifield) $\mathbb{Q}^+$ of positive rationals. Anyway, if S were such a (unitary) semiring with $1_S = 1_{\mathbb{Q}^+}$, then the additive semigroup $S(+)$ should be divisible. So far, all known examples of finitely generated additively divisible commutative semirings are additively idempotent. Hence a natural question arises, whether a finitely generated (commutative) semiring with divisible additive part has to be additively idempotent.

Analogous questions were studied for semigroups. According to [2, 2.5(iii)], there is a finitely generated non-commutative semigroup with a divisible element that is not idempotent. In this context it is of interest to ask whether there are non-finite but finitely generated divisible semigroups. This is in fact not true. Moreover, there exist infinite but finitely generated divisible (non-commutative) groups (see [3] and [8]).

The present short note initiates a study of (finitely generated) additively divisible commutative semirings. In particular, We show that an additively divisible commutative semiring is idempotent, provided that it is finitely generated and torsion. The one-generated case is treated in more detail and we prove that in the case that a one-generated additively divisible semirings possess no unit, it must contain an ideal of idempotent elements. However, even the one-generated case, in the mentioned conjecture, remains an open problem. Finally, we present a series of open questions about finitely generated commutative semirings and their equivalent versions.

1. Preliminaries

Throughout the paper, all algebraic structures involved (as semigroups, semirings, groups and rings) are assumed to be commutative, but, possibly, without additively and/or multiplicatively neutral elements. Consequently, a *semiring* is a

2010 *Mathematics Subject Classification.* 16Y60, 20M14.

Key words and phrases. commutative semiring, divisible semigroup, idempotent, torsion.

This work is a part of the research project MSM00210839 financed by MŠMT. The first author was supported by the Grant Agency of Czech Republic, #201/09/0296 and the second author by the project LC 505 of Eduard Čech's Center for Algebra and Geometry.

non-empty set equipped with two commutative and associative binary operations, an addition and a multiplication, such that the multiplication distributes over the addition. A semiring S is called a ring if the additive semigroup of S is a group and S is called a *parasemifield* if the multiplicative semigroup of S is a non-trivial group.

We will use the usual notation: $\mathbb{N}$ for the semiring of positive integers, $\mathbb{N}_0$ for the semiring of non-negative integers, $\mathbb{Q}^+$ for the parasemifield of positive rationals and $\mathbb{Q}$ for the field of rationals.

For all $k, t \in \mathbb{N}$ define a relation $\rho(k,t) = \{(m,n) \in \mathbb{N} \times \mathbb{N} |\ m - n \in \mathbb{Z}t \ \&\ (m \neq n \Rightarrow m, n \geq k)\}$. It is well known that the relations $\mathrm{id}_{\mathbb{N}}$ and $\rho(k,t)$, where $k, t \in \mathbb{N}$, are just all congruences of the additive semigroup $\mathbb{N}(+)$ (and of the semiring $\mathbb{N}$ as well).

Let S be a semiring. An element $a \in S$ is said to be *of finite order* if the cyclic subsemigroup $\mathbb{N}a = \{ka |\ k \in \mathbb{N}\}$, generated by the element a, is finite. We put $\mathrm{ord}(a) = \mathrm{card}(\mathbb{N}a)$ in this case. If $\mathcal{T}(S)$ denotes the set of elements of finite order, than either $\mathcal{T}(S) = \emptyset$ or $\mathcal{T}(S)$ is a *torsion* subsemiring (even an ideal) of S (i.e., every element of $\mathcal{T}(S)$ has finite order).

Lemma 1.1. *Let $\emptyset \neq A$ be subset of a semiring such that $\sup\{\mathrm{ord}(a) |\ a \in A\} < \infty$. Then $\sup\{\mathrm{ord}(b) |\ b \in \langle A \rangle\} < \infty$. Moreover, there is $r \in \mathbb{N}$ such that $2rb = rb$ for every $b \in \langle A \rangle$.*

Proof. Set $m = \sup\{\mathrm{ord}(a) |\ a \in A\} \in \mathbb{N}$. We have $\mathbb{N}a \cong \mathbb{N}(+)/\rho(k_a, t_a)$, where $k_a, t_a \in \mathbb{N}$, $k_a + t_a \leq m + 1$. Furthermore, $2m_a a = m_a a$ for some $m_a \in \mathbb{N}$, $m_a \leq m + 1$. Setting $r = (m+1)!$, we get $2rb = rb$ for every $b \in \langle A \rangle$. Of course, $\mathbb{N}b \cong \mathbb{N}(+)/\rho(k_b, t_b)$ and $\mathrm{ord}(b) = k_b + t_b - 1$. Since $2rb = rb$, we have $r \geq k_b$ and t_b divides $2r - r = r$. Consequently, $k_b + t_b - 1 \leq n$ for $n = 2r - 1 \in \mathbb{N}$. □

Lemma 1.2. *Let $a, b \in S$ be such that $ka = la + b$ for some $k, l \in \mathbb{N}$, $k \neq l$. If $\mathrm{ord}(b)$ is finite, then $\mathrm{ord}(a)$ is so.*

Proof. There are $m, n \in \mathbb{N}$ such that $m < n$ and $mb = nb$. Then $nka = nla + nb = nla + mb = (n-m)la + m(la + b) = (n-m)la + mka = ((n-m)l + mk)a$. Since $k \neq l$, we see that $(n-m)k \neq (n-m)l$ and $nk \neq (n-m)l + mk$. Consequently, $\mathrm{ord}(a)$ is finite. □

On a semiring S define a relation $\sigma_S = \{(a,b) \in S \times S |\ (\exists m \in \mathbb{N})\ ma = mb\}$. Clearly, σ_S is a congruence of S and $\sigma_{S/\sigma_S} = id$.

Lemma 1.3. *A semiring S is torsion, provided that the factor-semiring S/σ_S is torsion.*

Proof. For every $a \in S$ there are $k, l \in \mathbb{N}$ such that $(ka, la) \in \sigma_S$, $k < l$. Furthermore, there is $m \in \mathbb{N}$ with $mka = mla$. Clearly, $mk < ml$, and hence $\mathrm{ord}(a)$ is finite. □

Let S be a semiring and $o \notin S$ be a new element. Putting $x + o = o + x = x$ and $oo = xo = ox = o$ we get again a semiring $S \cup \{o\}$. Consider now the semiring $T = \mathbb{N}_0 \times (S \cup \{o\})$ equipped with component-wise addition and multiplication given as $(n,a)(m,b) = (nm, ma + nb + ab)$ for every $n, m \in \mathbb{N}_0$ and $a, b \in S \cup \{o\}$. Denote $\mathcal{U}(S) = T \setminus \{(0,o)\}$ the (unitary) subsemiring.

Now, S can always be treated as a natural unitary $\mathcal{U}(S)$-semimodule, with $(n,a)x = nx + ax$ for every $(n,a) \in \mathcal{U}(S)$, $x \in S$. (Here $(n,o)x = nx$ and $(0,a)x = ax$ for $n \in \mathbb{N}$ and $a, x \in S$.)

Lemma 1.4. *Let S be a semiring. If $w \in S$, $a, b, c \in \mathcal{U}(S)w$ and $m \in \mathbb{N}$ are such that $ma = mb$ and $mc = w$, then $a = b$.*

Proof. For every $d \in \mathcal{U}(S)w$, there is $\alpha_d \in \mathcal{U}(S)$ with $d = \alpha_d w$. Now, $a = \alpha_a w = \alpha_a mc = \alpha_c \alpha_a mw = \alpha_c ma = \alpha_c mb = \alpha_c \alpha_b mw = \alpha_b mc = \alpha_b w = b$. □

2. Additively divisible semirings

Let $S(= S(+))$ be a semigroup. An element $a \in S$ is called *divisible* (*uniquely divisible*, resp.) if for every $n \in \mathbb{N}$ there exists $b \in T$ (a unique, resp.) such that $a = nb$.

A semigroup S is called *divisible* (*uniquely divisible*, resp.) if every element of S is divisible (uniquely divisible, resp.). Clearly, S is divisible iff $S = nS$ for every $n \in \mathbb{N}$. The class of divisible semigroups is closed under taking homomorphic images and cartesian products and contains all divisible groups and all semilattices (i.e., idempotent semigroups).

A semiring is called *additively divisible* (*additively uniquely divisible*, resp.) if its additive part is a divisible semigroup (uniquely divisible semigroup, resp.). The semiring $\mathbb{Q}^+$ is additively (uniquely) divisible.

Note, that any semigroup $S(+)$ with an idempotent element e (i.e. $e + e = e$) can always be treated as a semiring with a constant multiplication given as $ab = e$ for every $a, b \in S$.

The following theorem is a consequence of [2, 2.5(i)].

Theorem 2.1. [2] *A semigroup S is finitely generated and divisible if and only if S is a finite semilattice.*

Theorem 2.2. *The following are equivalent for a commutative semiring S:*

(i) *S is additively idempotent.*
(ii) *S is additively divisible and bounded (i.e.* $\sup\{\mathrm{ord}(a) |\ a \in S\} < \infty$*).*
(iii) *S is additively uniquely divisible and torsion.*

Proof. (i)⇒(ii): Obvious.

(ii)⇒(i): By 1.1 there exists $n \in \mathbb{N}$ such that $2na = na$ for every $a \in S$. Since S is divisible, we have $a = nb$, and so $2a = 2nb = nb = a$.

(i)⇒(iii): Easy.

(iii)⇒(i): Since S is torsion, for every $a \in S$ there is $k \in \mathbb{N}$ such that $2ka = ka$. Now, S is additively uniquely divisible, hence $k(2a) = ka$ and $2a = a$. □

Corollary 2.3. *Let S be an divisible semiring. Then the factor-semiring S/σ_S is additively uniquely divisible.*

Moreover, if S is a torsion, then σ_S is just the smallest congruence of S such that the corresponding factor-semiring is additively idempotent.

Corollary 2.4. *Let S be an additively divisible semiring with $\mathcal{T}(S) \neq \emptyset$. Then $\mathcal{T}(S)$ is an additively divisible ideal in S.*

If, moreover, S is uniquely additively divisible, then $\mathcal{T}(S)$ is additively idempotent.

Example 2.5. Consider a non-trivial semilattice S and the quasicyclic p-group $\mathbb{Z}_{p^\infty}$. Then the product $S \times \mathbb{Z}_{p^\infty}$ is a torsion divisible semigroup that is neither a semilattice nor a group.

3. Additively divisible semirings - continued

Throughout this section, let S be an additively divisible semiring. The following assertion is easy to verify.

Proposition 3.1. *Assume that the semiring S is generated as an S-semimodule by a subset A such that* $\sup\{\mathrm{ord}(a) |\ a \in A\} < \infty$*. Then S is additively idempotent.*

Proof. Set $m = \sup\{\mathrm{ord}(a)|\ a \in A\} \in \mathbb{N}$ and put $B = \{b \in S|\ \mathrm{ord}(b) \leq m\}$. Then $A \subseteq B$ and $sb \in B$ for all $s \in S$ and $b \in B$. Consequently, $\langle B \rangle = S$ and, by 1.1, $\sup\{\mathrm{ord}(b)|\ b \in S\} < \infty$. Now, it remains to use 2.2. □

Corollary 3.2. *The semiring S is additively idempotent, provided that it is generated as an S-semimodule by a finite set of elements of finite orders.*

Corollary 3.3. *The semiring S is additively idempotent, provided that it is torsion and finitely generated.*

Remark 3.4. (i) The zero multiplication ring defined on $\mathbb{Z}_{p^\infty}$ is both additively divisible and additively torsion. Of course, the ring is neither additively idempotent nor finitely generated. The (semi)group $\mathbb{Z}_{p^\infty}(+)$ is not uniquely divisible.

(ii) Let R be a (non-zero) finitely generated ring (not necessary with unit). Then R has at least one maximal ideal I and the factor-ring R/I is a finitely generated simple ring. However, any such a ring is finite and consequently, R is not additively divisible.

Remark 3.5. Assume that S is additively uniquely divisible.

(i) For all $m, n \in \mathbb{N}$ and $a \in S$, there is a uniquely determined $b \in S$ such that $ma = nb$ and we put $(m/n)a = b$. If $m_1, n_1 \in \mathbb{N}$ and $b_1 \in S$ are such that $m/n = m_1/n_1$ and $m_1 a = n_1 b_1$, then $k = mn_1 = m_1 n$ and $kb_1 = mm_1 a = kb$ and $b_1 = b$. Consequently, we get a (scalar) multiplication $\mathbb{Q}^+ \times S \to S$ (one checks easily that $q(a_1 + a_2) = qa_1 + qa_2$, $(q_1 + q_2)a = q_1 a + q_2 a$, $q_1(q_2 a) = (q_1 q_2)a$ and $1a = a$ for all $q_1, q_2 \in \mathbb{Q}^+$ and $a_1, a_2, a \in S$) and S becomes a unitary $\mathbb{Q}^+$-semimodule. Furthermore, $qa_1 \cdot a_2 = a_1 \cdot qa_2$ for all $q \in \mathbb{Q}^+$ and $a_1, a_2 \in S$, and therefore S is a unitary $\mathbb{Q}^+$-algebra.

(ii) Let $a \in S$ be multiplicatively but not additively idempotent (i.e., $a^2 = a \neq 2a$). Put $Q = \mathbb{Q}^+ a$. Then Q is a subalgebra of the $\mathbb{Q}^+$-algebra S and the mapping $\varphi : q \mapsto qa$ is a homomorphism of the $\mathbb{Q}^+$-algebras and, of course, of the semirings as well. Since $a \neq 2a$, we have $\ker(\varphi) \neq \mathbb{Q}^+ \times \mathbb{Q}^+$. But $\mathbb{Q}^+$ is a congruence-simple semiring and it follows that $\ker(\varphi) = id$. Consequently, $Q \cong \mathbb{Q}^+$.

Put $T = Sa$. Then T is an ideal of the $\mathbb{Q}^+$-algebra S, $Q \subseteq T$ (we have $qa = qa \cdot a \in T$) and $a = 1_Q = 1_T$ is a multiplicatively neutral element of T. The mapping $s \mapsto sa$, $s \in S$, is a homomorphism of the $\mathbb{Q}^+$-algebras. Consequently, T is additively uniquely divisible. Furthermore, T is a finitely generated semiring, provided that S is so.

Proposition 3.6. *Assume that $1_S \in S$. Then:*

(i) *S is additively uniquely divisible.*
(ii) *Either S is additively idempotent or S contains a subsemiring Q such that $Q \cong \mathbb{Q}^+$ and $1_S = 1_Q$.*
(iii) *If $\mathrm{ord}(1_S)$ is finite, then S is additively idempotent.*

Proof. For every $m \in \mathbb{N}$, there is $s_m \in S$ such that $1_S = ms_m$. That is, $s_m = (m1_S)^{-1}$. If $ma = mb$, then $a = s_m ma = s_m mb = b$ and we see that S is additively uniquely divisible. The rest is clear from 3.5. □

Proposition 3.7. *If the semiring S is non-trivial and additively cancellative, then S is not finitely generated.*

Proof. The difference ring $R = S - S$ of S is additively divisible, and hence it is not finitely generated by 3.4(ii). Then S is not finitely generated either. □

4. One-generated additively divisible semirings

In this section, let S be an additively divisible semiring generated by a single element $w \in S$.

Proposition 4.1. *The semiring S is additively uniquely divisible.*

Proof. Follows from 1.4. □

Proposition 4.2. *The semiring S is additively idempotent, provided that* $\mathrm{ord}(w^m)$ *is finite for some $m \in \mathbb{N}$.*

Proof. Let $n \in \mathbb{N}$ be the smallest number with $\mathrm{ord}(w^n)$ finite. If $n = 1$, then the result follows from 3.2, and so we assume, for contrary, that $n \geq 2$. Since $S(+)$ is divisible, there are $v \in S$, $k \in \mathbb{N}_0$ and a polynomial $f(x) \in \mathbb{N}_0[x] \cdot x$ such that $w = 2v$ and $v = kw + w^2 f(w)$. Hence $w^{n-1} = 2kw^{n-1} + 2w^n f(w)$. By assumption, $2w^n f(w)$ is of finite order. If $k = 0$, then clearly $\mathrm{ord}(w^{n-1})$ is finite, and if $k \geq 1$, then $\mathrm{ord}(w^{n-1})$ is finite, by 1.2, the final contradiction. □

For $n \in \mathbb{N}$ denote $S^n = \langle \{a_1 \dots a_n |\ a_i \in S\} \rangle$.

Lemma 4.3. *If $u \in S$ is such that $w = wu$, then $u = 1_S$.*

Proof. Let $w = wu$. Since $S = \mathcal{U}(S)w$, for every $a \in S$ there is $\alpha_a \in \mathcal{U}(S)$ such that $a = \alpha_a w$. Hence $a = \alpha_a w = \alpha_a wu = au$ for every $a \in S$. Thus $u = 1_S$. □

Corollary 4.4. *$1_S \in S$ (i.e. S is unitary) if and only if $S^2 = S$. In this case:*

(i) *There exists w^{-1} in S.*
(ii) *$S^n = S^m$ for all $n, m \in \mathbb{N}$.*

Proof. Let $S^2 = S$. Then $w \in S = S^2$ and there is a non-zero polynomial $f(x) \in \mathbb{N}_0[x] \cdot x$ such that $w = wf(w)$. By 4.3, $f(w) = 1_S$. Now, since $1_S \in S = S^2$, we have similarly that $1_S = wu$ for some $u \in S$. Hence $w^{-1} = u \in S$. The rest is obvious. □

Lemma 4.5. *Let $k \in \mathbb{N}$, $k \geq 2$ and $u \in S \cup \{0\}$ be such that $w = kw + u$. Then there exists $a \in S \cup \{0\}$ such that $x = 2x + ax$ for every $x \in S$.*

Proof. Let $u = mw + wf(w)$, where $m \in \mathbb{N}_0$ and $f(x) \in \mathbb{N}_0[x] \cdot x$. If $f(x) = 0$, then $\mathrm{ord}(w)$ is finite, S is idempotent by 3.2 and we can put $a = 0$.

Hence assume that $f(x) \neq 0$. Put $n = m + k$ and $b = f(w) \in S$. We have $w = nw + wb$. Adding $(n-2)w$ to both sides of this equality, we get $(n-1)w = 2(n-1)w + wb$. Since w is a generator and S is additively divisible we have $b = (n-1)a$ for some $a \in S$ and for every $x \in S$ there is $\alpha_x \in \mathcal{U}(S)$ such that $x = (n-1)\alpha_x w$. Hence $x = \alpha_x(n-1)w = \alpha_x(2(n-1)w + (n-1)wa) = 2x + xa$ for every $x \in S$. □

Remark 4.6. Let $x, v \in S$ be such that $x = 2x + v$. Put $o = x + v$. Then:

(i) o is an idempotent and $x = x + o$.
(ii) The set $\{b \in S |\ x = x + b\}$ is a subsemigroup of $S(+)$ with o as an absorbing element.

Theorem 4.7. *If $1_S \notin S$, then there exists $a \in S \cup \{0\}$ such that $\{u + au |\ u \in S\}$ is an ideal in S consisting of idempotent elements.*

Proof. Since S is divisible, there is $f \in \mathbb{N}_0[x]$ such that $w = 2f(w)$. By 4.3, we get that $f(x) = kx + xg(x)$ for some $k \in \mathbb{N}$, $k \geq 2$ and $g \in \mathbb{N}_0[x]$. Now, by 4.5 there is $a \in S \cup \{0\}$ such that $u = 2u + au$ for every $u \in S$. Finally, $u + au$ is idempotent for every $u \in S$ by 4.6. □

Clearly, S is additively divisible, iff for every $n \in \mathbb{N}$ there is a non-zero polynomial $f_n(x) \in \mathbb{N}_0[x] \cdot x$ such that $w = nf_n(w)$. By [2, 2.5(i)], every divisible element in a finitely generated commutative semigroup has to be idempotent. Assuming now, that the degree of the polynomials f_n is bounded, we get that S is

additively idempotent. To illustrate some other situations and techniques see the next examples.

Example 4.8. Let $n, m, k, l \in \mathbb{N}$ be such that $n^{l-1} \neq m^{k-1}$ and suppose that $w = nw^k$ and $w = mw^l$. Then S is additively idempotent.

Indeed, $nm^k w^{kl} = n(mw^l)^k = nw^k = w = mw^l = m(nw^k)^l = mn^l w^{kl}$. Since $nm^k \neq mn^l$, w^{kl} is of finite order and S is additively idempotent by 4.2.

Example 4.9. Let $w = 2(w + w^2)$ and $w = 3(w + w^3)$. Then S is additively idempotent.

Indeed, adding these two equalities we get $2w = 5w + 2w^2 + 3w^3$. Now, since $w^2 = 2w^2 + 2w^3$ we can substitute $2w = 5w + (2w^2 + 2w^3) + w^3 = 5w + w^2 + w^3$. Hence $4w = 10w + (2w^2 + 2w^3)$ and by the same substitution we get $4w = 10w + w^2$. Finally, $8w = 20w + 2w^2 = 18w + (2w + 2w^2) = 19w$ and w is of finite order. Thus S is additively idempotent by 4.2.

Even when a one-generated semiring does not have a unit, we can use another one that will contain a unit and will share some properties with the original semiring (see 4.10).

Proposition 4.10. *$T_S = \{\varphi \in \mathrm{End}(S(+)) |\ (\exists \alpha \in \mathcal{U}(S))(\forall x \in S)\ \varphi(x) = \alpha x\}$ is an additively divisible commutative semiring that has a unit id_S and is generated by the set $\{\mathrm{id}_S, \varphi_w\}$, where $\varphi_w(x) = wx$.*

Moreover, S is additively idempotent if and only if T_S is so.

Proof. We only need to show, that $\mathrm{id}_S \in T_S$ is an additively divisible element. Let $n \in \mathbb{N}$. Then $w = na$ for some $a \in S$ and $a = \alpha_a w$ for some $\alpha_a \in \mathcal{U}(S)$. Now, for every $x \in S$ there is $\alpha_x \in \mathcal{U}(S)$ such that $x = \alpha_x w$. Hence $x = \alpha_x w = \alpha_x n \alpha_a w = n\alpha_a x$. Thus $\mathrm{id}_S = n\alpha_a$. The rest is easy. □

5. A few conjectures

In this last section we present some other open questions that are influenced by our main problem (namely by the conjecture (A) - see below).

Example 5.1. Given a multiplicative abelian group G and an element $o \notin G$, put $U(G) = G \cup \{o\}$ and define addition and multiplication on $U(G)$ (extending the multiplication on G) by $x + y = xo = ox = o$ for all $x, y \in U(G)$. Then $U(G)$ becomes an ideal-simple semiring.

Consider the following statements:

- (A) Every finitely generated additively divisible semiring is additively idempotent.
- (A1) Every finitely generated additively uniquely divisible semiring is additively idempotent.
- (B) No finitely generated semiring contains a copy of $\mathbb{Q}^+$.
- (B1) No finitely generated semiring with a unit element contains a copy of $\mathbb{Q}^+$ sharing the unit.
- (B2) Every finitely generated additively divisible semiring with a unit is additively idempotent.
- (C) Every parasemifield, that is finitely generated as a semiring, is additively idempotent.
- (C1) Every finitely generated infinite and ideal-simple semiring is additively idempotent or a copy of the semiring $U(G)$ (see 5.1) for an infinite finitely generated abelian group G.

Proposition 5.2. (A) $\Leftrightarrow$ (A1) $\Rightarrow$ (B) $\Leftrightarrow$ (B1) $\Leftrightarrow$ (B2) $\Rightarrow$ (C) $\Leftrightarrow$ (C1).

Proof. First, it is clear that (A) $\Rightarrow$ (A1), (B) $\Rightarrow$ (B1) and (A) $\Rightarrow$ (B2). Furthermore, (C) $\Leftrightarrow$ (C1) by [5, 5.1]. Now, assume that (A1) is true and let S be a finitely generated additively divisible semiring. By 2.3, S/σ_S is additively uniquely divisible and, of course, this semiring inherits the property of being finitely generated. By (A1), the semiring S/σ_S is additively idempotent, and hence the semiring S is additively torsion by 1.3. Finally, S is additively idempotent by 3.3. We have shown that (A1) $\Rightarrow$ (A) and consequently, (A) $\Leftrightarrow$ (A1).

Next, let (B1) be true and let S be a finitely generated semiring containing a subsemiring $Q \cong \mathbb{Q}^+$. Put $P = S \cdot 1_Q$. Then P is an ideal of S, $1_Q = 1_P$, $Q \subseteq P$ and the map $s \mapsto s1_Q$ is a homomorphism of S onto P. Thus P is a finitely generated semiring and this is a contradiction with (B1). We have shown that (B1) $\Rightarrow$ (B) and consequently, (B) $\Leftrightarrow$ (B1).

The implication (B2) $\Rightarrow$ (B1) is easy, since every semiring S with a unit element 1_S that contains a subsemiring $Q \cong \mathbb{Q}^+$ with $1_S = 1_{\mathbb{Q}^+}$ is additively divisible (for $a \in S$ and $m \in \mathbb{N}$ choose $b = (m1_S)^{-1}a \in S$ and get $a = mb$).

The implication (B) $\Rightarrow$ (B2) follows immediately from 3.6.

We have shown that (B) $\Rightarrow$ (B2) $\Rightarrow$ (B1) $\Leftrightarrow$ (B).

Finally, the implication (B2) $\Rightarrow$ (C) follows from the fact that every parasemifield S is additively divisible, since the prime parasemifield of S containg 1_S is isomorhic either to $\mathbb{Q}^+$ or to the trivial semiring. □

The conjecture (C) was confirmed for the one-generated case in [6] and for the two-generated case in [4].

Note that using the Birkhoff's theorem we can consider an equivalent version of the conjecture (A):

(A') Every finitely generated subdirectly irreducible additively divisible semiring is additively idempotent.

Of course, it would be sufficient if such a semiring were finite. Unfortunately, this is not true. Assume for instance the semiring $S = V(G)$ with $G = \mathbb{Z}(+)$. By [1, 10.1], S is simple and two-generated. Nevertheless, it is an open question whether also one-generated subdirectly irreducible additively divisible semiring can be infinite.

Finally, Mal'cev [7] proved that every finitely generated commutative semigroup is residually finite (i.e. it is a subdirect product of finite semigroups). Notice, that also the additive part of a free finitely generated additively idempotent semiring is a residually finite semigroup. If this is true also for every finitely generated additively divisible semiring, we get a nice positive answer to the conjecture (A).

References

[1] R. El Bashir, J. Hurt, A. Jančařík, T. Kepka, *Simple commutative semirings*, J. Algebra **236** (2001), 277–306.

[2] W. E. Clark, W. C. Holland, G. J. Székely, *Decompositions in discrete semigroups*, Studia Sc. Math. Hungarica **34** (1998), 15–23.

[3] V. S. Guba, *A finitely generated complete group*, Izv. Akad. Nauk SSSR, Ser. Mat. **50(5)** (1986), 883-924 (English translation: Math. USSR, Izv. **29** (1987), 233–277).

[4] J. Ježek, V. Kala, T. Kepka, *Finitely generated algebraic structures with various divisibility conditions*, Forum Math. **24(2)** (2012), 379–397.

[5] V. Kala, T. Kepka, *A note on finitely generated ideal-simple commutative semirings*, Comment. Math. Univ. Carolinae **49** (2008), 1–9.

[6] V. Kala, T. Kepka, M. Korbelář, *Notes on commutative parasemifields*, Comment. Math. Univ. Carolinae **50(4)** (2009), 521–533.

[7] A. I. Mal'cev, *On homomorphisms into finite groups*, Lecture Notes of Ivanovsk Pedagog. Inst. **18** (1958), 49–60.

[8] A. Yu. Olshanskii, *Geometry of Defining Relations in Groups*, Nauka, Moscow, (1989) (in Russian) (English translation: Kluwer, 1991).

Charles University, Faculty of Mathematics and Physics, Department of Algebra, Sokolovská 83, 186 75 Prague 8, Czech Republic
E-mail address: kepka@karlin.mff.cuni.cz

Department of Mathematics and Statistics, Faculty of Science, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic
E-mail address: miroslav.korbelar@gmail.com